# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY 36852

DATE: JUN 13 EST

Request for Confirmation of a Ceiling Increase; a Twelve-Month SUBJECT: Exemption; and Confirmation of the Initial Verbal Approval for

Funding the Removal Action at the White Bridge Road Site, Meyersville, Morris County, New Jersey - ACTION MEMORANDUM

FROM:

Michael Neill, On-Scene Coordinator  $_{\text{TO}}$ : Removal Action Branch

Constantine Sidamon-Eristoff Regional Administrator

Kathleen C. Callahan, Director Emergency and Remedial Response Division

Site I. D. - 6G
Category of Removal - Emergency
National Significance - Yes (asbestos primary contaminant)

### I. PURPOSE

Friable and free asbestos in large quantities have been found on the White Bridge Road National Priority List (NPL) site, a satellite site to the Asbestos Dump, NPL site. During an August 1990 site inspection, the Environmental Protection Agency (EPA) site evaluation team observed white dust being kicked up by horses exercising on a training oval. Subsequent sampling from the oval revealed 5% asbestos present. Further sampling from the on-site dwelling showed asbestos present in dust samples. As a result of these findings, the Agency for Toxic Substances and Disease Registry (ATSDR) verbally issued a preliminary health consultation, followed by a memorandum, advising EPA to temporarily relocate residents at the White Bridge Road site. Subsequently, an addendum to ATSDR's April 1989 Health Assessment was issued. ATSDR issued a preliminary Public Health Advisory in September 1990, and a final Public Health Advisory on December 20, 1990 for this site. ATSDR has indicated that steps should be taken to reduce the exposure to the asbestos dust at the sites, perform further sampling, and dissociate humans from free asbestos at the site. The Removal Site Evaluation and the ATSDR documents are attached.

On August 23, 1990, verbal approval was given, by the Region II Director of the Emergency and Remedial Response Division (ERRD), to initiate activities necessary to protect all persons from exposure to asbestos at the site. Concurrence was also received from Headquarters since asbestos is the primary contaminant at the site. The initial Project Ceiling was established at \$249,000 with \$200,000 in mitigation contracting and \$49,000 set aside for TAT and EPA costs. On September 13, 1990, verbal approval was given by William Muszynski, Deputy Regional Administrator, to raise the project ceiling to \$750,000 to complete initial removal actions. The authority for this approval comes from the Comprehensive Environmental Response,

100 GB

On August 23, 1990, verbal approval was given, by the Region II Director of the Emergency and Remedial Response Division (ERRD), to initiate activities necessary to protect all persons from exposure to asbestos at the site. Concurrence was also received from Headquarters since asbestos is the primary contaminant at the site. The initial Project Ceiling was established at \$249,000 with \$200,000 in mitigation contracting and \$49,000 set aside for TAT and EPA costs. On September 13, 1990, verbal approval was given by William Muszynski, Deputy Regional Administrator, to raise the project ceiling to \$750,000 to complete initial removal actions. The authority for this approval comes from the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA), as delegated to the EPA regional offices.

By this memorandum, these approvals are confirmed. The increase in funds was necessary since the extent of contamination to be addressed was found to be considerably larger than originally estimated. In addition, TAT and EPA costs increased since the owner/residents declined to be relocated at the time of the removal action and therefore, special consideration and planning was required. The new mitigation contracting amount is \$310,000; TAT cost is \$220,000; and EPA cost is \$220,000. The new project ceiling is \$750,000.

Since uncertainty exists when all actions can be considered complete, and other actions may have to be taken at different times over the project period, an exemption to the twelve-month time limitation is also requested.

### II. SITE CONDITIONS AND BACKGROUND

The Asbestos Dump Superfund site is an NPL site comprised of four subsites. The "main" site is known as the Millington site and is currently being remediated as Operable Unit 1. There are three "satellite" sites known as the New Vernon Road site, the White Bridge Road site and the Deitzman Tract. The New Vernon Road and the White Bridge Road sites comprise Operable Unit 2. The Deitzman Tract comprises Operable Unit 3. Removal actions, one of which is the subject of this memorandum, were initiated in 1990 at the New Vernon Road and White Bridge Road sites.

The White Bridge Road site is located at 651 White Bridge Road, Meyersville, Morris County, New Jersey. It consists of approximately 12 acres and is bounded by New Vernon Road, White Bridge Road and the Great Swamp National Wildlife Refuge. There is one residence located on site, as well as horse stables and a riding track and grazing areas for horses. There are two adult residents living on-site as well as 18 horses reportedly boarding in the stables. The residents operate a horse boarding business on-site. There are other private residences to the north and west of the site (see Figure 1).

areas suspected of having been impacted by the asbestos waste materials from the Millington facility, including the White Bridge Road site.

Subsequent to the issuance of the Order, it was deemed appropriate to split remedial activities into operable units. The Millington site is Operable Unit 1. EPA has selected a remedy for this portion of the site which is set forth in a September 1988 Record of Decision. Operable Unit 2 of the site is comprised of the two satellite sites which are residential properties; the New Vernon Road and the White Bridge Road sites. The third satellite site, known as the Deitzman Tract, comprises Operable Unit 3 of the site.

In April 1989, ATSDR issued a Health Assessment for the site which recommended, among other things, that additional sampling be performed. In August 1990, based on ATSDR's recommendations, and as part of the NPL site assessment effort, the Removal Action Branch collected and analyzed soil and dust samples. The sampling program was performed with the Environmental Response Team (ERT). The site was found to be potentially unsafe for residents, the owners/boarders/trainers of the horses and nearby residents. The sampling results showed 5% asbestos on the training ring and 2% asbestos in the entrance lane to the ring. Due to the high levels of asbestos, EPA determined that an immediate removal action was necessary to address the imminent threat posed by the site. The removal action was initiated based on verbal authorization by the Director of ERRD.

The removal action was initiated in August 1990. The removal activities performed to date include: collecting and analyzing samples from the site; covering primary areas of asbestos contamination, including the horse riding ring and lane leading to the ring, with geotextile fabric to restrict access and reduce the potential for airborne releases; and, erecting signs and temporary fencing to restrict access to areas with surface contamination.

The removal action served to stabilize and temporarily reduce the risks posed by exposure to asbestos by site residents and others. During the fall of 1990, concurrent with the removal action, EPA initiated a remedial investigation (RI) at the site. This RI included extensive soil and air sampling to characterize site contamination. To minimize the disturbance of asbestos, the RI sampling was performed prior to the placement of the geotextile fabric over areas of asbestos contamination.

The data collected during the RI has characterized the nature and extent of asbestos contamination at the site, and is currently serving as a basis for selection of a final remedy. This remedy will be documented in a subsequent Record of Decision.

### B. Quantities/Types of Substances Present

The White Bridge Road site contains asbestos waste in the vicinity of the horse riding ring and the lane leading to the ring, as well as one area on the west-central portion of the site. The thickness of the asbestos waste ranges from 0 to 12 feet. An estimated 21,324 cubic yards of asbestos waste is present on the site.

Samples collected on August 2, 1990 from the horse riding ring and entrance lane to the ring were analyzed by Transmission Electron Microscopy (TEM), and found to contain 5% and 2% chrysotile asbestos, respectively. A dust sample collected from the dwelling on August 24, 1990 indicated the presence of free chrysotile asbestos at a concentration of less than 1%.

Asbestos, when it is friable, is designated as a CERCLA hazardous substance under 40 CFR Table 302.2. The term "friable" means that the material can easily be crumbled by hand, and therefore, is likely to emit fibers into the air.

### C. Site Conditions

Asbestos has been found in various surface and subsurface areas on the site. Weathering has made the material friable and the action of horses on the riding ring and vehicles using the entrance lane, had pulverized the tile chips into dust, resulting in a major health threat.

### III. THREAT

An April 1989 ATSDR Health Assessment stated that since the potential exists for the transport of appreciable levels of asbestos into the air, the potential exists for significant exposure to asbestos at concentrations of immediate public health concern. An August 29, 1990 ATSDR memorandum to the Region states that an immediate and significant health threat exists and actions should be taken to abate the threat. An addendum to the original health assessment, confirming the health threat, was issued on September 14, 1990. A Health Advisory, based on the addendum, was issued by ATSDR on December 20, 1990. This Health Advisory contained a number of recommendations including that: affected residents be dissociated from asbestos fibers in indoor air; additional sampling be performed; and access to asbestos contaminated areas be restricted.

Since the probability exists for asbestos fibers to become airborne and continue to do so unless mitigative measures are taken, a major health threat exists at this site.

The most significant pathway by which airborne asbestos affects humans is through the respiratory system. Asbestos exposure may

cause two primary classes of health effects. The first is asbestosis, a non-malignant disease characterized by a progressive scarring of the lungs and pleura. This condition progresses very slowly over many years, and may continue even after the exposure has ceased. As microscopic scarring builds up, the lungs become stiff and restrict breathing. The other major class of asbestos-related health effect is mesothelioma, a type of lung cancer, which may occur even after low level exposure to the asbestos. All asbestos-related malignancies have a latency period. There usually is a considerable time interval between the exposure and when the adverse effects are seen. This latency period may vary from ten to forty years, although some cases have been documented where the effects were seen soon after exposure.

### IV. ENDANGERMENT DETERMINATION

Actual or threatened releases of hazardous substances from this site, if not addressed by the implementation the response action selected in this Action Memorandum, presented an imminent and substantial endangerment to public health or welfare.

### V. EXEMPTION FROM STATUTORY LIMITS

The White Bridge Road site meets the following criteria, as prescribed in CERCLA Section 104(c)(1)(C), which is required to exceed the twelve-month time limit for removal actions:

The continued response actions are otherwise appropriate and consistent with the remedial actions to be taken.

Extensive coordination and close monitoring of activities with the Remedial Program and the Office of Regional Counsel has ensured that the public has been protected during the removal action. Actions taken were consistent and appropriate with long term solutions to this immediate health threat. Removal activities were coordinated with RI sampling.

In addition, the response actions implemented are appropriate under Section 104(c)(1)(C) of CERCLA, as amended by the Superfund Amendments and Reauthorization Act (SARA), since the following criteria have been met:

### A. Consistency

Asbestos abatement measures at this site consisted of posting signs and installing fencing, and temporarily covering the exposed asbestos areas with geotextile fabric. All activities were closely coordinated with the Remedial Program and consideration was given to the final remedial action. Actions undertaken to date are similar to those at like sites in other regions.

### B. Appropriateness

ATSDR issued a Health Advisory on December 20, 1990. The Health Advisory for this site calls for:

- dissociating humans from free/friable asbestos;
- additional sampling;
- posting warning signs;
- restricting access to known areas of asbestos contamination; and,
- eliminating activities which could cause the asbestos to become airborne.

The removal action accomplished these goals in an expeditious manner, including reducing the possibility of further off-site migration through runoff or erosion caused by air motion. The measures taken during the removal action have stabilized the site through temporary measures. The geotextile fabric which covers the major areas of asbestos contamination, will be maintained until a permanent remedy is constructed at the site.

### VI. REMOVAL ACTIVITIES

The removal activities performed to date include: collecting and analyzing samples from the site; covering primary areas of asbestos contamination, including the horse riding ring and lane leading to the ring with geotextile fabric to restrict access and reduce the potential for airborne releases; and, erecting signs and temporary fencing to restrict access to areas with surface contamination.

### VII. SUMMARY OF COSTS

Summary of Costs	Current Ceiling	Proposed Ceiling
Extramural Costs ERCS IAG ERT TAT	\$ 200,000  \$ 30,000	\$ 310,000   \$ 220,000
Total Extramural Costs	<b>\$ 230,0</b> 00	\$ 530,000
Total Intramural Costs	<b>\$ 19,0</b> 00	\$ 220,000
PROJECT CEILING	<b>\$ 249,0</b> 00	\$ 750,000

### VIII. EXPECTED CHANGE IN THE SITUATION SHOULD ACTION BE DELAYED OR NOT TAKEN

The proposed actions were taken based, in part, on the ATSDR's Health Advisory. Had the actions not been taken, the residents and any person entering the site, or coming near the site, could

1

have potentially been exposed to free asbestos fibers and been at risk of diseases caused by exposure to these fibers.

### IX. ENFORCEMENT

National Gypsum Company has been identified as a PRP for the Asbestos Dump site. An Order was issued in 1985 to National Gypsum for the performance of RI/FS activities at the site, including the three satellite sites. In 1987, National Gypsum submitted an RI/FS report which documented RI/FS activities at the sites. At that time, EPA determined that adequate information was presented to select a final remedy for the Millington portion of the site, but that additional investigation was necessary to fully characterize the three satellite sites, including the White Bridge Road site, prior to selection of final remedies.


Subsequently, upon determining that a substantial threat existed at the New Vernon Road and White Bridge Road sites based on 1990 sampling, EPA performed the removal action described above.

### X. RECOMMENDATIONS

Conditions at the White Bridge Road site continue to meet the criteria for a CERCLA removal action under 40 CFR 300.415, in that a release of a hazardous substance to the environment has occurred and could continue to occur. Qualifying criteria include the following:

- (b)(2)(i) Actual or potential exposure to nearby human populations, animals, or the food chain from hazardous substances or pollutants or contaminants;
- (b)(2)(iv) High levels of hazardous substances or pollutants or contaminants in soils largely at or near the surface, that may migrate;
- (b)(2)(v) Weather conditions that may cause hazardous substances or pollutants or contaminants to migrate or be released;
- (b)(2)(vii) The availability of other appropriate federal or state response mechanisms to respond to the release;
- (b)(2)(viii) Other situations or factors that may pose threats to public health welfare or the environment.

Therefore, I request your approval to confirm the original project ceiling and the new project ceiling. The new project ceiling is \$750,000. Of this amount, \$310,000 is for mitigation contracting, and \$440,000 is for TAT and EPA costs.



## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY REGION II

DATE: SEP 11 1990

JBJECT: Removal Site Evaluation for Whitebridge Road Asbestos Millington Dump Site (NPL), Meyersville, New Jersey

FROM: Nick Magriples, On-Scene Coordinator Hid Magriples
Removal Action Branch

TO:

The File

### I. INTRODUCTION

As part of the National Priorities List (NPL) Removal Assessments, the United States Environmental Protection Agency's (U.S. EPA) Removal Action Branch (RAB) has reviewed conditions at the Whitebridge Road Asbestos Millington Site (Whitebridge Road) for Comprehensive Environmental Response, Compensation and Liability Act (CERCLA) Removal Action consideration.

The Whitebridge Road Site is a satellite site, of operable unit two, under the NPL project - Asbestos Millington Dump site. The other two satellite sites are the Dietzman Tract and the New Vernon Road site. Operable unit one is the Millington site.

A release of asbestos to the soil, through past disposal and filling operations, has occurred at the Whitebridge Road Site. Both the dirt roadway on the eastern edge of the property, and the horse riding track to which it leads, are covered/paved with pulverized asbestos tile that becomes dusty during dry periods. Additionally, there are areas of asbestos fill on the southeastern portion of the property and at the edge of the grazing field. The dirt roadway appears to be accessed by both horses and vehicles. The latter appears to be used for moving the horse hurdles. The probability of asbestos fibers being disturbed by the wind or contact, and becoming airborne, appears to be high.

The Agency of Toxic Substances and Disease Registry has verbally issued a preliminary commentary on the health threat, stating that the Whitebridge Road Site poses an imminent health threat, and calling for; medical monitoring of the on-site residents, and additional sampling.

Based on recent analytical data revealing elevated levels of fibrous asbestos (2-5% chrysotile) in the surface soils, the potential for a serious release of asbestos to the air exists at this site. Additionally, a lower level of fibrous asbestos (< 1% chrysotile) was detected in a dust sample collected from the house on the site. Due to the presence of residents, at the house on the site, horse riders that use the track and board their horses in stables at the site, and the potential for off-site migration, a removal action should be conducted to mitigate

787

ļ

the potential and existing threats that are posed by the presence of asbestos in the surface soils.

Region 2 Removal Ranking System: 9

### II. PERSONNEL INVOLVED

The following U.S. EPA personnel were directly involved in the Removal Assessment conducted for the White Bridge Road Site: Nick Magriples (201-906-6930) and John Witkowski (201-321-6739) of the Removal Action Branch, Edison, New Jersey.

### III. SITE SETTING

The Whitebridge Road Site is located in Meyersville, New Jersey and consists of approximately 12 acres of land off New Vernon Road and bounded by the Great Swamp National Wildlife Refuge to the east and south, and private residences to the north and west (see Figure 1). The address is 651 Whitebridge Road. There are two residents living on-site and 18 horses reportedly boarding in stables at the site. There are five other residences on Whitebridge Road (between New Vernon Road and the Great Swamp), within approximately 700 feet of the site. One of them is directly across the street.

### IV. BACKGROUND

The land at the site was used as a farm from 1945 until 1969. From 1970 until 1975, landfilling operations by the National Gypsum Corporation were performed. The refuse included asbestos tiles and siding. Following termination of the landfilling, the owner converted the property into a horse farm with stables, a riding ring, and pasture fields.

According to the draft Remedial Investigation (RI) report, the Whitebridge Road site has asbestos contamination in the eastern section of the property and along the main driveway. The main landfill area consists of the area south of, and including, the riding track and a small portion of the grazing field. The depth of the asbestos wastes averages between two to four and one-half feet. Towards the southeastern corner of the site it is reportedly at least 10 feet deep.

The Whitebridge Road Site is a satellite site of the National Priorities List (NPL) Asbestos Millington Dump site. A 106 Order issued by the U.S. EPA on April 4, 1985 required National Gypsum to perform a Remedial Investigation/Feasibility Study (RI/FS) on the entire Asbestos Dump. Subsequent to issuance of the Order, it was deemed appropriate to split the work into two operable units. The RI for operable unit two has been conducted (1987), however it has not been approved by the U.S. EPA Remedial Program. Further activities have not been conducted for operable

unit two.

In August 1989, the U.S. EPA issued an Interagency Agreement (IAG) with the U.S. Fish and Wildlife Service (U.S. FWS) to provide technical support for the RI/FS for operable unit two.

### V. <u>SITE ACTIVITIES/OBSERVATIONS</u>

The RAB conducted a reconnaissance of the site on March 22, 1990 and a site visit, with the U.S. FWS project manager, Sid Mitra and representatives of the U.S. EPA Emergency Response Team (ERT), on June 28, 1990. The purpose of the site visit was to determine what additional sampling would be required, if any, to complete the removal assessment for the site. ERT was present, at the OSC's request, to provide technical assistance for the evaluation.

Based on ERT recommendations, the RAB collected soil and dust samples from the site on August 2, 1990. The soil samples were collected from various points at the surface of the dirt roadway, the riding track, and the edge of the grazing field. Samples were analyzed for % asbestos content and type of asbestos fiber using the Transmission Electron Microscopy Method (TEM) by International Asbestos Testing Laboratories, Maple Shade, New Jersey (see Materials section below for an explanation of the results). On August 24, 1990, ERT collected a dust sample from within the house on the site and analyzed it for both Phased Light Microscopy (PLM) and TEM.

The site consists of a two story building where the owners reside, a garage, and several sheds and stables that are used for the horses (see Figure 2). The roadway on the northwest side of the site, leading to all of the above structures, is paved with asphalt. There is a large, grazing field for the horses in the center of the site which takes up the majority of the property. A pond, approximately 100 feet in diameter, is situated in the northern portion of the grazing field. Trees line the property, east of the home, along Whitebridge Road. The riding track is approximately 250 feet long by 125 feet wide and is situated approximately 350 feet from the house and stables. The dirt roadway is approximately 250 feet long.

The asbestos appears to be most exposed and friable on the riding track, where it has been pulverized and mixed with dirt by the horses. It should be noted that during the August 2 sampling event, when the riding track was very dry due to the lack of rain, the material was easily made airborne with ordinary walking.

The dirt roadway consists of small asbestos chips that have been worn and crushed from repeated usage. The eastern portion of the grazing field adjacent to the riding track also contains

pulverized asbestos mixed in with the soil. Further west into the field, chips are visible at the base of the crabgrass. Along the southern end of the riding track and towards the south larger pieces (6-12 inches) of asbestos tile are visible at the surface.

### VI. MATERIALS ON-SITE

Samples collected by the RAB on August 2, 1990 confirm visual observations that friable asbestos is present throughout the driveway of the Whitebridge Road Site (see Figure 1 for approximate sample locations). Samples BL008 from the dirt roadway indicates 2% chrysotile content. Samples BL009 - B012 from the riding track and adjoining areas indicate 5% chrysotile content.

A sample collected by ERT on August 24, 1990 revealed fibrous asbestos (< 1% chrysotile) in the dust from within the house on the site.

Asbestos is designated as a CERCLA hazardous substance under 40 CFR Table 302.4, when it is friable. Friable means that it can be crumbled with hand pressure, and therefore, is likely to emit fibers when disturbed.

### VII. THREAT

A release of asbestos to the soil, through past disposal and filling operations, has occurred at the Whitebridge Road Site. Both the dirt roadway on the eastern edge of the property, and the horse riding track to which it leads, are covered/paved with pulverized asbestos tile that becomes dusty during dry periods. Additionally, there are areas of asbestos fill on the southeastern portion of the property and at the edge of the grazing field. The dirt roadway appears to be accessed by both horses and vehicles. The latter appears to be used for moving the horse hurdles. The probability of asbestos fibers being disturbed by the wind or from contact, and becoming airborne, appears to be high. Since the RI has not been approved to date, remediation of the site is not scheduled for the near future. Due to the presence of this contaminant source, and its friability, the potential for a release of asbestos to the air in the future will continue to exist and is likely to occur.

An April 1989 ATSDR Health Assessment stated that since the potential exists for the transport of appreciable levels of asbestos, the potential exists for significant exposure to asbestos at concentrations of long-term public health concern.

ATSDR has verbally issued a preliminary commentary on the health threat at the Whitebridge Road Site, stating that the site poses an imminent health threat based on the levels of fibrous asbestos present in the surface soils (2-5% chrysotile). It called for

medical monitoring of the on-site residents, and additional sampling.

Asbestos is a general term used to describe minerals that tend to form fibers when they are broken. These minerals are formed under conditions of very high heat and pressure deep in within the earth, and they are resistant to the types of temperatures and pressures found in our environment at the surface. Because their chemical composition is unchangeable, an asbestos mineral will always break into fibers. Large fibers have the potential to break into smaller ones, which eventually results in its reduction to microscopic size. Because of their small size, shape and lightness, these fibers act are more like a gas than a dust.

The most important human exposure pathway for asbestos is the inhalation of respirable asbestos fibers. The ingestion of fibers may also be an exposure pathway of concern for workers or children who may come into contact with site materials. In addition to environmental exposures, the improper handling of work clothing from on-site workers may also pose a danger. Workers can carry the fibers home in their clothing and hair, and expose other family members.

Asbestos exposure may cause two primary classes of health effects. The first is asbestosis, a non-malignant disease characterized by a progressive scarring of the lung and pleura. This condition progresses slowly over many decades, and may continue even after the asbestos exposure has ceased. As microscopic scarring builds up, the lungs become stiff and restricted with thickening in the walls of the breathing spaces. The stiffening of the lungs, when severe, can make it difficult to breathe.

The other major class of asbestos-related health effects is mesothelioma and lung cancer after apparently trivial exposure to asbestos.

All asbestos-related malignancies have a latency period. There is a considerable time interval between asbestos exposure and when lung cancer, mesothelioma, or the other asbestos-related cancers are seen. This latency period may vary from twenty to forty years, although some cases may occur earlier.

### VIII. CONCLUSIONS

A release of asbestos to the soil, through past disposal and filling operations, has occurred at the Whitebridge Road Site. Both the dirt roadway on the eastern edge of the property, and the horse riding track to which it leads, are covered/paved with pulverized asbestos tile that becomes dusty during dry periods. Additionally, there are areas of asbestos fill on the

southeastern portion of the property and at the edge of the grazing field. The dirt roadway appears to be accessed by both horses and vehicles.

The probability of asbestos fibers being disturbed by the wind or contact, and becoming airborne, appears to be high. Based on this, ATSDR's preliminary commentary, the recent analytical data revealing elevated levels of fibrous asbestos in the surface soils (2-5% chrysotile) and in the house (< 1% chrysotile), the presence of residents at the house on the site, and horse riders that use the track and board their horses in stables at the site, a removal action should be conducted to mitigate the potential and existing threats that are posed by the presence of asbestos. Based on the final ATSDR Health Advisory, an addendum to this RSE may be required.

### IX. RECOMMENDATIONS

Conditions at the Whitebridge Road Site meet the criteria for a CERCLA removal action under 40 Code of Federal Regulations (CFR) 300.400 and the National Oil and Hazardous Substances Pollution Contingency Plan. Qualifying criteria include the following:

- 40 CFR 300.415(b)(2)(i) Actual or potential exposure to nearby human populations, animals or the food chain from hazardous substances or pollutants or contaminants;
- 40 CFR 300.415(b)(2)(iv) High levels of hazardous substances or pollutants or contaminants in soils largely at or near the surface, that may migrate;
- 40 CFR 300.415(b)(2)(v) Weather conditions that may cause hazardous substances or pollutants or contaminants to migrate or be released;
- 40 CFR 300.415(b)(2)(vii) The availability of other appropriate federal or state response mechanisms to respond to the release;
- 40 CFR 300.415(b)(2)(viii) Other situations or factors that may pose threats to public health or welfare or the environment.

The following mitigative measures, at a minimum, are recommended for the Whitebridge Road Site:


- all roads should be paved or covered in some manner;
- the riding track should be covered in some manner, and restored to a rideable condition;
- the areas of the grazing field that contain exposed


asbestos should either be excavated and/or covered;

- the area south of the riding track that contains exposed asbestos should either be excavated and/or covered;
- the grassy area of the property that the homeowner regularly cuts should be investigated further with sampling and a determination made on the need for any mitigation;
- further sampling and analysis may be warranted to establish any additional or more extensive mitigative measures.

Although remedial alternatives have not been established, the removal action should consider, to the extent possible, potential long term remedial actions when developing the scope of work. Although the conditions at the site appear to have been in existence for some time, an expedited time critical response is appropriate for the Whitebridge Road Site based on the exigency of the circumstances at the site.

- cc. G. Zachos
  - J. Witkowski
  - M. Ferriola
  - M. Neill





ABD 001 2091

John

FAX Dick SAI KIE

FINAL DRAFT

\*\*00 NOT CITE OR QUOTE\*\*

comments due to

DRAFT

AGENCY FOR TOXIC BUBSTANCES AND DISEASE REGISTRY
DIVISION OF HEALTH ASSESSMENT AND CONSULTATION
PUBLIC HEALTH ADVISORY

ASBESTOS DISPOSAL SITES NPL SITE NEW VERNON ROAD SUBSITE AND WHITE BRIDGE ROAD SUBSITE

PASSAIC TOWNSHIP, MORRIS COUNTY, NEW JERSEY

September 21, 1990

INTRODUCTION

As a result of a recommendation in a Health Assessment of the Aubestos Disposal Sites, the U.S. Environmental Protection Agency (U.S. EPA) conducted soil and dust sampling for asbestos at the Asbestos Disposal Sites near Meyersville, New Jersey. Based on the analytical results of that sampling, the Agency for Toxic Substances and Disease Registry (ATSDR) has determined that the New Vernon Road Site and the White Bridge Road Site represent an imminent and substantial threat to public health. The source of this threat is the asbestos contamination that has been found in the soil and in the homes located on these two sites. The risk of exposure to asbestos is increased for the residents in any home contaminated with free asbestos fibers. Asbestos is a known human carcinogen with a prolonged latency period of between ten and thirty years between exposure and the onset of disease. Advarse health effects are known to occur after an exposure of limited duration. Health effects are known to occur after inhalation of asbestos fibers and may occur through ingestion

BRAFT HEALTH ADVISORY ASSESSES DISPOSAL SILES

DRAFT

-----

AR OF: Reptminer 21, 1990 ORIGINAL DATE: August 30, 1990

## FINAL DRAFT

of asbestos fibers.

## DRAFT

Potential health effects include: asbestosis, a physical injury of the lung tissue caused by the asbestos fibers; lung cancer; mesotheliomas, tumors formed within the thin membrane surrounding internal organs, primarily caused by exposure to asbestos; and, gastrointestinal cancer, including the colon and esophagus.

Because of the known carcinogenicity of asbestos and the likelihood of exposure at these sites, the ATSDR recommends that: residents in houses known to be contaminated with asbestos be dissociated from the contamination existing as free fibers; dwellings adjacent to these two sites be sampled for free asbestos fibers and, if found, the residents be dissociated from the contomination; activities at the sites which tend to increase sirborne particulates be restricted to areas where asbestos contomination is known not to exist; the homes of workers at these sites be sampled for free asbestos fibers and, if found, the workers and their families be dissociated from the contemination.

Attached Comments and the contemination.

The purposes of this public health advisory are to notify the U.S. EPA, the New Jersey State Department of Health, and the public that an imminish and substantial human health threat exists at these sites, and to bring to their attention the ATSDR's concerns and recommendations for the protection of the public health.

2

DRAFT HEALTH ADVISORY ASSESTED DISPUSAL SITES AT OFF Exptudier 21, 1999 ORIGINAL BATE: AUGUST 30, 1990

### DRAP "DO NOT CITE OR QUOTE"

## DRAFT

**あ**ひなひょうりじょうつ

### BACKGROUND

The Agency for Toxic Substances and Disease Registry (ATSDR) received a request from the U.S. EPA for an avaluation of the health threat posed by asbestos contamination in the soll and in one dwalling at the New Varnon Road Site. The contemination was found through a sampling event recommended in the Health Assessment (HA), prepared by the ATSDR, deted April 10, 1989. The sampling was conducted in July 1990 at two of the subsites associated with this site: New Vornon Road Site and White Bridge Road Site.

Analysis of the samples revealed a maximum concentration of 5% by volume chrysotile asbestos in the soil and 2% by volume in a residence. All of the twelve samples collected contained at least 24 by volume chrysotile.

The New Vernon Road Sire consists of approximately 30 acres of land off New Vernon Road in Meyersville, New Jersey in Passaic Township. In the late 1960s, asbestos refusa from an ashestus processing plant in Millington was landfilled on the site at two separate locations. These locations are now called the filled pond area and the main landfill area. The refuse consisted of loose asbestos fibers, broken asbestos tilos, and broken asbestos siding.

DRAFT HEALTH ADVISORY ASSESTED DISPOSAL SITES

AS IFE Teplestur 21, 1550 CRICINAL PAIR: ACRUME 30, 1990

DRAFT

### DRAFT

## FINAL DRAFT

The White Bridge Road Site consists of approximately 12 acres in Meyersville. The site is now a horse farm. From 1970 to 1975, weetes similar to those disposed at the New Vernon Road site were landfilled in the eastern portion of the site in and around what has become a riding track.

Both the White Bridge Road Site and the New Varnon Road Site are located in a primarily rural area. A combined total of 15-20 off-site residences are potentially impacted by any migration of the wastes from the two sites. Additional site descriptive information and demographics can be found in the NA.

Asbestos is a group of six naturally occurring fibrous minerals. Asbentos is a known human carcinogen and is one of the primary causes of mesothelioms. Mesotheliomas are tumors srising from the thin membrane surrounding internal organs. Inhelation of asbestos fibers may lead to fibrotic lung disease (asbestosis), cancer of the lung, the pleurs, and the peritoneum. There is some evidence that inhelation and ingestion of asbestos fibers may lead to an increased risk of gastrointestinal cancer.

There is a substantial latency period of between ten and thirty years between the exposure and the occurrence of apparent health effects. Some human and animal studies have indicated that adverse effects may occur

DRAFT HEALTH ADVISORY AGREGIUS DISPOSAL SITES AS UF: Replanter 21, 1990 ORIGINAL DATE: AUGAST 50, 1990

DRAFT

## DRAFT

### TINAL DRAFT

after exposures of limited duration. In order for exposure to occur, the asbastos must exist as free fibers capable of becoming airborns.

The length and diameter of fiber is important in determining the ultimate effect of the exposure. For instance, fibers less than 0.5 micrometers in diameter are most active in producing tumors. However, there is little doubt, based on human epidemiological studies and animal studies, that all types of asbestos, including chrysotile, can cause cancer.

All concentrations of ashostos fibers studied to date have domonstrated an excess cancer risk, as reported in the National Institute for Occupational Safety and Health (NIOSH) Revised Recommended Asbestos Standard. A marked enhancement of the risk of lung carcinoms in exposed workers or populations who also smoke eigerettes has been noted in human epidemiology studies. The ATSDR Toxicological Profile for Asbestos indicates this increased risk may be as high as ten times the nonsmoker risk.

### BASIS FOR ADVISORY

There is ample opportunity for human exposure to the chrysotile asbestos at both of these subsites. The respective owners spend a great deal of time in the outdoors, either at the stables or working with equipment, and children play outdoors on site.

5

DRAFT HEALTH ADVISORY ASSESTED DISPUSAL SITES

AS OF: September 21, 1990 URIGINAL DATE: August 30, 1990

DRAFT

ABD 001 20

-F.740+1011111

## DRAFT

### FINAL DRAFT \*\*\*po NOT CITE OR CLOTE\*\*

DRAFT

Contamination may be brought into the home as well. There have been documented incidents in which asbestos workers have carried contamination home on their clothing and on their person. In this situation, it is possible that children and pets which frequent the site may also carry contamination into the home.

The high concentration of asbestos in the soil outside the homes represents a serious threat to the occupant's health. Samples taken in the dwellings on site clearly establish the presence of free asbestos fibers in the homes. Continued exposure to free asbestos fibers, regardless of the concentration, represents an imminent and substantial health threat to exposed individuals.

### CONCLUSIONS

Residents on site in the homes with free asbestos fibers face an imminint and substantial health threat from exposure to asbestos. Workers at the New Vernon Road Site and workers and customers at the horse farm and riding stables at the White Bridge Road Site also face an increased risk of doveloping adverse health effects. The familles of these workers and customers may also be at risk.

DRAFE HEALTH ADVISING AGGERICE DISPOSAL SILES

AE OF: Emitender 21, 1990 ORIGINAL DATE: August 30, 1990

DRAFT

## FINAL DRAFT

## DRAFT

### RECOMMENDATIONS AND PROPOSED ACTIONS

TO-NO O'D ELM CD LEN'LENCH . ..

The ATSDR, in consultation with the New Jersey State Department of Health, will develop criteria, based on exposure, for selection of residents and make available a medical monitoring program to those selected residents.

The ATSDR recommends the following actions he taken to mitigate this health threat:

- The U.S. EPA should dissociate those residents with free ambestos fibers in the home from the contamination.
- 2. Additional sampling should be performed by the U.S. EFA to determine if homes adjacent to these sites contain free subsetos fibers caused by off-site migration.
- 3. Additional sampling should be performed by the U.S. EPA in the homes of workers and customers of the businesses located at these sites.
- 4. The U.S. EPA should post warning signs in the vicinity of the horse track at the White Bridge Road Site to advise nustomars of the hazards at the site.

DRAFT HEALTH AUVISORY ASSESTOS DISPOSAL SITES AR OF: Repterbar 21, 1990 ORIGINAL DATE: August 30, 1990

## DRAFT

## FINAL DRAFT \*\*BO NOT CLTE ON GLOTE\*\*

- 5. The U.S. EPA or the property owners should restrict access,
  authorized or unauthorized, to those areas known to be contaminated
  with asbestos.
- 6. The U.S. EPA or the property owners should eliminate activities which tend to increase airborne particulates in those areas known to be contaminated with asbestos.

For additional information, please contact the ATSDR at the following address:

Robert C. Williams, P.E.

Director, Division of Health Assessment and Consultation
Agency for Toxic Substances and Disease Registry
1600 Clifton Road, NE, MS E-32
Atlanta, Georgia 30333
(404) 639-0610
FTS 236-0610

DRAFT HEALTH ADVISORY ASSESTOR DISPUSAL BITES AS OF: September 21, 1990 ORIGINAL DATE: August 30, 1990



### DEPARTMENT OF HEALTH & HUMAN SERVICES

Public Health Service Centers for Disease Control

Memorandum

"September 14, 1990 Date

From

Environmental Health Scientist, Emergency Response and Consultation Branch, Division of Health Assessment and Consultation, ATSDR (232)

Subject

Addendum Asbestos Disposal Sites NPL Site Millington (Morris County), New Jersey

To

Arthur Block Public Health Advisor ATEDR Regional Services Region II

Through: Director, DEAC, ATEDR (E32) Chief, RPB, DHAC, ATEDR (232)

Chief, ERCB, DEAC, ATSDR (E32) E



Attached is an addendum to the Asbestos Disposal Sites NFL Site Health Assessment. The addendum is based on a review of the analytical results of soil and residential dust sampling. These samples were collected by the Environmental Protection Agency (EPA) in July 1990 in response to a recommendation contained in the ATSDR Health Assessment. The results indicated that soil at two subsites contained high concentrations of chrysotile asbestos (5% by volume) and at least one dvelling was contaminated by high levels of chrysotile asbestos (2% by volume). The two subsites addressed in this addendum are the New Vernon Road Site and the White Bridge Road Site in Passaid Township, New Jersey. The addendum is not applicable to any other subsite associated with this RPL site.

Flease contact me if you have any questions regarding this addendum

### Asbestos Disposal Sites RPL Site Morris County, Millington, New Jersey CERCLIS NO. NJD980654149 Addendum to the Easith Assessment

### BACKGROUND AND STATEMENT OF ISSUES

The Agency for Toxic Substance and Disease Registry (ATSDR) received a request from the U.S. Environmental Protection Agency (EPA) for an evaluation of the health threat posed by asbestos contamination in the soil and in residences at the Asbestos Disposal Sites BPL Site. The contamination was found as a result of sampling recommended in the Health Assessment (HA), prepared by ATSDR, dated April 10, 1989. Twelve samples were collected in July 1990 at two of the subsites associated with this site: New Vernon Road Site and White Bridge Road Site.

Analysis of the samples revesled a maximum concentration of 5% by volume chrysotile asbestos in the soil and 2% by volume in a residence. The residential sample was collected from the occupant's vacuum cleaner bag. All of the twelve samples collected contained at least 2% by volume chrysotile. No other type of asbestos fibers were reported at these subsites.

On the New Vernon Road Bits seven samples were collected. The samples were collected in one of two dwellings, in a small garage, along a dirt road leading between the two buildings to the main disposal area, and slong a dirt path bordering the disposal area. At the White Bridge Road Site, five samples were collected. These sample were collected on or near the riding track and the dirt path leading up to the track.

The New Vernon Road Site consists of approximately 30 acres of land off New Vernon Road in Meyersville, New Jersey, in Fassaic Township. In the late 1960s, asbestos refuse from an asbestos processing plant in Millington was landfilled on the site at two separate locations. locations are now called the filled pond area and the main landfill area. The refuse consisted of loose asbestos fibers, broken asbestos tiles, and broken asbestos miding. Previous investigations detected asbestos waste in the pond ares in front of the residence, the main landfill, the dirt road, and in the vicinity of a shed near the residence. The property owner, his wife and two children, and one adult tenent live on the site. The owner employs approximately 3 other parsons in his on-site business. These workers spend part of each workday on site.

ABD

The White Bridge Road Site consists of approximately 12 acres in Neyersville. The site is now a horse farm. From 1970 to 1975, waster similar to those disposed of at the New Vernon Road site were landfilled in the eastern portion of the site in and around what has become a riding track. Asbestos contemination has been found on the riding track, the dirt path leading to the riding track, and in the adjacent grazing field. The property owner and his wife live on the site. The horse farm and it's associated riding track is a commercial enterprise employing approximately three individuals besides the owner. These employees work on site.

Both the White Bridge Road Site and the New Vernon Road Site are located in a primarily rural area. A combined total of 15-20 off-site residences are potentially impacted by off-site migration of the wastes. Additional site descriptive information and demographics can be found in the KA.

Follow-up sampling by the Environmental Protection Agency in August 1990 found loose fibers of chrysotile asbestos in the dust in several residences with distreters ranging from 0.02 um to 0.2 um. The concentration of asbestos in the dust was reported as less than 1% by volume in all samples. These dust samples were collected in areas where contamination was expected to be found as recommended by the ATSDR. These data have not undergons quality assurance review and should be considered preliminary.

### DOCUMENTS AND INFORMATION REVIEWED

Health Assessment for Asbestos Disposal Site. Prepared by Agency for . Toxic Substances and Disease Registry. April 10, 1989.

Revised Recommended Asbestos Standard. Prepared by National Institute for Occupational Safety and Health. December 1976.

Draft Toxicological Profile for Asbastos. Prepared by Agency for Toxic Substances and Disease Registry. February 16, 1990.

Excerpts from Sampling Report: New Vernon Road Site and White Bridge Road Site, Asbestos Dump Sites. Prepared by Fred C. Hart Associates, Inc., Subcontractor to Roy F. Weston, Inc. Prepared for: U.S. Environmental Protection Agency, Region II. Undated.

Conference Call between Agency for Toxic Substances and Disease Registry, Division of Health Assessment and Consultation, Emergency Response and Consultation Branch; Public Health Advisor, Agency for Toxic Substances and Disease Registry, Region II; and, New Jersey Department of Health. August 21, 1990. Refer to ERCB Superfund Record of Communications dated August 27, 1990.

Conversation between Hr. Richard Nickle, Agency for Toxic Substances and Disease Registry and Hr. William Howard, Centers for Disease Control. August 30, 1990. Refer to ATSDR Superfund Record of Communications dated August 31, 1990.

Conversation between Mr. Richard Hickle, Agency for Toxic Substances and Disease Registry and Mr. Raymond McQueen, Rational Asbestos Council. August 30, 1990. Refer to ATSDR Superfund Record of Communications dated August 31, 1990.

TO

Facsimile transmission of preliminary analytical data. Transmitted by: Mr. Arthur Block, Public Health Advisor, Regional Services, Region II, Agency for Toxic Substances and Disease Registry, New York. Transmitted to: Mr. Richard Bickle, Division of Health Assessment and Consultation, Emergency Response and Consultation Branch, Agency for Toxic Substances and Disease Registry. August 5, 1990

### DISCUSSION

Asbestos is a group of six naturally occurring fibrous minerals. Asbestos is a known human carcinogen and is one of the primary causes of mesothelioms. Mesotheliomas are tumors arising from the thin membranes surrounding internal organs. Inhalation of asbestos fibers may lead to fibrotic lung disease (asbestosis), cancer of the lung, the pleurs, and the peritoneum. There is some evidence that inhalation and ingestion of asbestos fibers may lead to an increased risk of gastrointestinal cancer. In order for exposure to occur, the asbestos must exist as free fibers capable of becoming airborne. There is a substantial latency period of between ten and thirty years between the time exposure first occurs and the occurrence of apparent health effects. Some human and animal studies have indicated that adverse effects may occur even after relatively short periods of exposures.

The length and diameter of fiber is important in determining the ultimate effect of exposure. However, there is little doubt, based on human epidemiological studies and animal studies, that all types of asbestos, including chrysotile, can cause cancer. Chrysotile asbestos was the only form of asbestos fiber identified at these subsites. Diameters of the fibers found were in the range of 0.02 to 0.2 um. Fibers less than 0.5 um in diameter are most active in producing tumors (NIOSH Recommended Standard). Inhalation of chrysotile fibers can also lead to asbestosis, a chronic lung disorder.

All concentrations of asbestos fibers studied to date demonstrated an excess cancer risk, as reported in the NIOSH Recommended Standard. Both the NIOSH Recommended Standard and the ATEDR Toxicological Profile report a marked enhancement of the risk of lung carcinoms in exposed workers or populations who also smoke eigerettes. The Toxicological Profile indicates this increased risk may be as high as ten times the nonsmoker risk.

In making conclusions based on data obtained from vacuum cleaner bag samples, the following factors should be considered:

ABD 001 2

. ; .

11.14.60

- o The concentration of asbestos in the vacuum bag may reflect higher or lower concentrations than is actually present in the home.
- O Although primarily used in the household, many vacuum cleaners are used elsewhere (e.g., cleaning the interior of a car). This introduces other sources of contamination which may skew the enalytical results.
- O During vacuuming, some fibers may pass through the bag and be exhausted into the ambient air of the home. Thus, the action of the vacuum may increase the potential for exposure by increasing the number of fibers in the air in the breathing some.

There is, however, ample opportunity for human exposure to the chrysotile asbestos to occur at both of these subsites. The respective owners spend a great deal of time in the outdoors, either at the stables or working with equipment, and children play outdoors on site. Contamination may be brought into the home as well. There have been documented incidents in which asbestos workers have carried contamination home on their clothing and on their person. In this situation, it is possible that children and pets which frequent the site may also carry contamination into the home.

### - CONCLUSIONS

As described previously, excessive risk has been demonstrated at all fiber concentrations studied as reported by NIOSH in their recommended standard. The high concentration of chrysotile asbestos in the soil outside the homes represent a serious threat to the occupant's health. Therefore, ATSDR concludes as follows:

- 1. Based on the information available, the residents at both subsites face an imminent and substantial health threat from exposura to asbestos through inhalation and possibly ingestion. Smokers would face an increased risk of asbestos related health effects as compared to non-smokers.
- 2. A potential health threst also exists for the surrounding population if the friable asbestos on-site migrates off-site.
- 3. A potential health threat also exists for workers at the New Vernon Road Site and workers and customers at the horse farm and riding stables at the White Bridge Road Site. The families of these workers and customers may be at risk as well.

### RECOYMENDATIONS

1. The exposure of the residents on these sites to asbestos should be terminated.

ABD 001

- 4 -

- 2. The occupants of the dwelling in which the asbastos has been found should undergo medical examination. Feriodic examinations after an initial baseline is established should be considered by the attending physician since asbastos exposure has a substantial latency period.
- 3. Additional sampling in adjacent homes should be undertaken. If dust samples are collected, they should be from relatively inaccessible areas, such as the top of bookshelves and under refrigerators.
- 4. The data collected during the additional sampling recommended should be evaluated to determine if further action is necessary. ATSDR would be available to perform this evaluation.
- 5. Access to the sites should be restricted.
- 6. Activities which tend to generate dust, including horseback riding, should be terminated in areas known to be contaminated with asbestos.
- 7. Warning signs specifying the exposure risk should be posted in the vicinity of the horse track at the White Bridge Road Bits. The signs should remain in place until mitigative efforts alleviate the health threat.

ABD 001

- 3 -

### Preparers of Report

Emergency Response Reviewer

Richard A. Mickle
Emergency Response Coordinator
Emergency Response and
Consultation Branch
Division of Health Assessment
and Consultation

Toxicological Reviewer

Allen Sustan
Senior Toxicologist
Emergency Response and
Consultation Branch
Division of Health Assessment
and Consultation

- 6 -

2105



Memorandum

Date August 29, 1990

From Arthur Block D ATSDR Regional Representative

Subject Asbestos Dump Site: Morris County, New Jersey ATSDR Health Review and Comment

Ms. Kathleen Callahan
Deputy Director, ERRD, EPA Region II

On August 22, 1990 the Environmental Protection Agency (EPA), Region II, requested that the Agency for Toxic Substances and Disease Registry (ATSDR) Region II, review and comment on residential sampling data obtained from a National Priority List (NPL) site located in Morris County, New Jersey, named Asbestos Dump Site (ADS). EPA's Field Investigation Team (F.I.T) had taken soil and household samples in and around two of A.D.S. operable units (o.u.); the Mt. Vernon Road Site (MVR) and the White Eridge Road Site (WBR).

The F.I.T. investigation discovered elevated levels of chrysotile asbestos (5%) in driveways and soil samples in and around the residential roads and pathways of both the MVR and WBR Sites. Additionally, 5% levels of asbestos was found in the horse track area of the WBR Site and a 2% level of chrysotile asbestos was found in a dust sample taken from the Tielman Household's vacuum cleaner located in the MVR Site.

### Recommendations:

In consultation and agreement with the Emergency Response Branch (ERB) of ATSDR, ATSDR, Region II recommends the following:

- (1) ATSDR concludes that at the reported levels of asbestos found, that we recommend A.D.S. an <u>Immediate and Significant Health Threat</u>.
- (2) ATSDR further recommends that the occupants of the Tielman household located adjacent to the MVR Site be Immediately and Temporarily Relocated.
- (3) ATSDR recommends that the residential houses in the immediate vicinity of the MVR and WBR Sites be immediately sampled for the presence of asbestos. Results to be forwarded to ATSDR for review and comment.

ABD 001 2106

Kathleen Callahan Deputy Director

August 29, 1990

### Follow-up:

On April 10, 1989 ATSDR released a Health Assessment on the Asbestos Dump Site. ATSDR will prepare an addendum to this Health Assessment stating the above recommendations and will shortly release a Preliminary Health Advisory on this issue.

Please direct any questions or comments of this review to Arthur Block (212-264-7662) or Lisa Voyce (264-7662), Regional Representatives, ATSDR, Region II.

cc: Constantine Sidamon-Eristoff
William J. Muszynski
Richard Caspe
Richard Salkie
Sharon Jaffers
Nick Margripolis
Mike Ferriola
Pat Seppi
George Buynoski
Richard Nickle
Suzanne Simon
Jonathan Savrin
Jim Pasqualo
Gregory Ulirsch

ABD 001

001 210

DEP SAULIE

CS-E MUSTANK DAMUSUI Caope Sinu

Agency for Toxic Substances and Disease Registry Atlanta GA 30333

DEC 2 0 1990

Mr. Constantine Sidamon-Eristoff
Regional Administrator
U.S. Environmental Protection Agency
Region II
Jacob K. Javitts Federal Building
New York, New York 10278

Dear Mr. Sidamon-Eristoff:

This letter is in reference to an enclosed Public Health Advisory for current and potential exposures to hazardous wastes, specifically chrysotile asbestos, at the New Vernon Road Site and the White Bridge Road Site in Meyersville, Passaic Township, New Jersey. These two sites are subsites of the Asbestos Disposal Sites NPL Site in Millington, New Jersey.

The Agency for Toxic Substances and Disease Registry (ATSDR) reviewed the analytical results of air, soil, and residential dust sampling at these sites. These samples were collected by the U.S. Environmental Protection Agency (EPA) beginning in August 1990. The initial results indicated that soil at two subsites contained high concentrations of chrysotile asbestos (5 percent by volume) and at least one dwelling was contaminated by high levels of chrysotile asbestos (2 percent by volume). The Advisory is not applicable to any other subsite associated with this NPL site.

The enclosed Public Health Advisory expresses our concerns and addresses measures to mitigate the risk to human health. By separate letter, Dr. William L. Roper, ATSDR Administrator, has notified the EPA Administrator of this Advisory.

Sincerely yours,

Barry D. Johnson Ph.D. Assistant Surgeon General

Assistant Administrator

Enclosure

BD 001 210

# AGENCY FOR TOXIC SUBSTANCES AND DISEASE REGISTRY DIVISION OF HEALTH ASSESSMENT AND CONSULTATION PUBLIC HEALTH ADVISORY

### ASBESTOS DISPOSAL SITES NPL SITE NEW VERNON ROAD SUBSITE AND WEITE BRIDGE ROAD SUBSITE

### PASSAIC TOWNSHIP, MORRIS COUNTY, NEW JERSEY

December 20, 1990

### INTRODUCTION

The U.S. Environmental Protection Agency (EPA), Region II, conducted soil and dust sampling for asbestos at subsites of the Asbestos Disposal Sites in Meyersville, New Jersey, as part of a program designed to assess the need for removal actions at NPL Sites. Based on the analytical results of that sampling, the Agency for Toxic Substances and Disease Registry (ATSDR) has determined that contamination at the New Vernon Road Site and the White Bridge Road Site presents a public health concern. This Public Health Advisory is issued to notify the EPA, the New Jersey State Department of Health, and the public that the presence of asbestos at these two subsites represents an imminent and substantial threat to human health.

The health threat results from chrysotile asbestos contamination in the soil and in the homes located on these two sites. The risk of exposure to free asbestos fibers is increased for the residents of any home in the general area which is contaminated with site-related asbestos. Persons who work at or visit these sites may also be at increased health risk because of the potential for exposure to free asbestos fibers at concentrations above background. Background implies levels of asbestos detected in similar rural areas remote from the source of site-related asbestos.

Chrysotile, the substance of health concern in these homes, is one of a group of naturally occurring fibrous silicate minerals, generally referred to as asbestos. The ATSDR considers the different mineral forms of asbestos to be known human carcinogens with a prolonged latency period of between 10 and 30 years between exposure and the onset of disease. Adverse health effects can occur after an exposure of limited duration. Health effects are known to occur after inhalation of asbestos fibers and may occur through ingestion of asbestos fibers. [2,3]

Potential health effects include: asbestosis (a physical injury of the lung tissue caused by the asbestos fibers); lung cancer; mesotheliomas (malignant tumors formed within the thin membrane surrounding internal organs, primarily caused by exposure to asbestos); and, gastrointestinal cancer, including the colon and esophagus. [3]

Because of the known carcinogenicity of asbestos and the likelihood of exposure at these sites, ATSDR recommendations include: (1) residents in on-site homes known to be contaminated with free asbestos fibers be dissociated from the contamination existing as free fibers; (2) buildings, including the dwellings, adjacent to these two sites be sampled for free asbestos fibers, and, if fibers are found at

comparable concentrations to the on-site dwellings, the occupants be dissociated from the contamination; (3) activities at the sites that would increase airborne particulates be restricted at areas where asbestos contamination is known to exist; (4) the homes of employees of the businesses at these sites be sampled for free asbestos fibers, and, if fibers are found at concentrations comparable to the on-site dwelling, the workers and their families be dissociated from the contamination. The EPA Region II is aggressively implementing these recommendations and has substantially reduced the concentrations of the asbestos fibers in one of the dwellings at the New Vernon Road Site.

The purposes of this Public Health Advisory are to notify the EPA, the New Jersey State Department of Health, and the public of the substantial human health hazard at these sites, and to bring to their attention ATSDR's concerns and recommendations for the protection of the public health.

### BACKGROUND

The ATSDR received a request from EPA Region II for an evaluation of the health hazard posed by asbestos contamination in the soil and in one dwelling at the New Vernon Road Site. The contamination was found through a sampling event conducted as part of a removal assessment program of NPL sites. The sampling was conducted by EPA Region II in August 1990 at two of the subsites associated with this site: New Vernon Road Site and White Bridge Road Site.

Analysis of the samples, according to an analytical method for determining bulk asbestos content, revealed a maximum concentration of 5 percent by volume chrysotile asbestos in the soil and 2 percent by volume in a residential vacuum cleaner bag. All 12 samples collected contained at least 2 percent by volume chrysotile. In making conclusions based on data obtained from vacuum cleaner bag samples, the following factors should be considered:

- The concentration of asbestos in the vacuum bag may reflect higher or lower concentrations than are actually present in the home.
- Although primarily used in the household, many vacuum cleaners are used elsewhere (e.g., cleaning the interior of a car). This introduces other sources of contamination which may skew the analytical results.
- During vacuuming, some fibers may pass through the bag and be exhausted into the ambient air of the home. Thus, vacuuming may increase the potential for exposure by increasing the number of fibers in the air in the breathing zone.

In September 1990, sampling of household dust by EPA Region II in the homes on-site and in some adjacent homes confirmed the presence of free asbestos fibers. At least one sample from each home contained detectable asbestos fibers. The levels detected were below quantification levels for the analytical methods used.

\BD 001 21

Indoor air samples collected in October 1990 allowed for a comparison of the levels of asbestos found in the home at the New Vernon Road Site to levels of asbestos present in control homes, which would be considered background concentrations. At the recommendation of ATSDR, EPA selected two homes which were representative of homes in the area of the subsites, but known to be unassociated with site-related asbestos (control homes). These two homes were sampled as was the dwelling on the New Vernon Road Site. Preliminary results indicate that no asbestos particulates or fibers were present in the control homes. The samples from the New Vernon Road Site contained a total concentration of asbestos fibers of 3000 fibers per cubic meter (f/m³) [0.0003 fibers per cubic centimeter (f/cc)]. Characterization of the New Vernon Road samples showed that the concentration of fibers over 5 micrometers (um) in length was 0.0013 f/cc or 1300 f/m³. [11]

Following this round of air sampling, a removal action was completed in the dwelling and the air was resampled. Results indicated a concentration of  $1900 \text{ f/m}^3$ . The samples from the control homes were collected using a passive technique (i.e., normal household activities) while the air sample in the dwelling was collected using an aggressive technique (e.g., fans or blowers agitated the dust and fibers). The control samples are, therefore, indicative of normal exposures, while the dwelling sample is indicative of worst case exposures. [10]

The New Vernon Road Site consists of approximately 30 acres of land and two dwellings off New Vernon Road in Meyersville, New Jersey, in Passaic Township. In the late 1960s, asbestos refuse from an asbestos processing plant in Millington was placed in landfills on the site at two separate locations. These locations are now called the filled pond area and the main landfill area. The refuse consisted of loose asbestos fibers, broken asbestos tiles, and broken asbestos siding.

The White Bridge Road Site consists of approximately 12 acres and one dwelling in Meyersville. The site is now a horse farm. From 1970 to 1975, wastes similar to those disposed at the New Vernon Road Site were placed in a landfill in the eastern portion of the site in and around what has become a riding track.

Both the White Bridge Road Site and the New Vernon Road Site are located in a primarily rural area. A combined total of 15-20 off-site residences are potentially impacted by any migration of the wastes from the two sites. This number of potentially-impacted residences is based on the observations of the ATSDR Regional Representative over the course of several site visits and includes the homes of the employees of the businesses on these sites. Additional site descriptive information and demographics can be found in the Health Assessment. [1]

On the New Vernon Road Site, the property owner operates a tree surgery business which is reported to employ between three and four persons. These employees may be exposed to the asbestos on-site in the course of their employment. This exposure is likely to be less than 2 hours per day since most of the employees' tasks are off-site. On the White Bridge Road Site, a stable and riding track for horses is operated by the property owner. Two to three employees and, to a

lesser extent, an unknown number of customers may be exposed to asbestos on-site while grooming and handling the horses, especially in the area of the riding track.

Chrysotile is one of a group of six naturally occurring fibrous silicate minerals generally referred to as asbestos. Asbestos is a known human carcinogen and is one of the primary causes of mesothelioma. Mesotheliomas are tumors arising from the thin membrane surrounding internal organs. Inhalation of asbestos fibers can lead to fibrotic lung disease (asbestosis), cancer of the lung, the pleura, and the peritoneum. There is some evidence that inhalation and ingestion of asbestos fibers may lead to an increased risk of gastrointestinal cancer. However, Chrysotile has been shown to cause all of the adverse health effects associated with asbestos exposure. [2,3]

There is a substantial latency period of between 10 and 30 years between the exposure and the occurrence of apparent adverse health effects. Some human and animal studies have indicated that adverse health effects can occur after exposures of limited duration. In order for exposure to occur, the asbestos must exist as free fibers capable of becoming airborne. [2,3]

The length and diameter of fiber is important in determining the ultimate effect of the exposure. For instance, fibers less than 0.5 micrometers in diameter are those most active in producing tumors. [2] The ATSDR considers all mineral forms of asbestos, including chrysotile, to be a hazard to human health, based on human epidemiological studies and animal studies.

To date, exposure to all concentrations of asbestos fibers have demonstrated an excess cancer risk. [2,3] A marked enhancement of the risk of lung carcinoma in exposed workers or populations who also smoke cigarettes has been noted in human epidemiology studies. [2,3] The ATSDR Toxicological Profile for Asbestos indicates the increased risk associated with smoking may be as high as ten times the risk for nonsmokers.

Ambient concentrations of asbestos in urban areas have been reported to be less than 100 nanograms total asbestos per cubic meter of air (ng/m³). [9] In one study described on page 75 of NIOSH's document, Revised Recommended Asbestos Standard, the concentration of asbestos in a building insulated with asbestos averaged 6,000 fibers of chrysotile per cubic meter of air. [2; Nicholson, Rohl, and Weisman, 1975] In another study completed later and described on page 82 of ATSDR's Draft Toxicological Profile for Asbestos, asbestos in indoor air was reported in the range of 20 to 6,000 fibers of asbestos per cubic meter. Ambient air concentration in rural settings across the country range from 1 to 3 orders of magnitude smaller than indoor air. [3; Nicholson, 1987] The ATSDR considers that a mass of 1 nanogram of asbestos may contain a sufficient number of fibers to create a health threat.

#### BASIS FOR ADVISORY

There is ample opportunity for human exposure to chrysotile asbestos at the New Vernon Road Site and the White Bridge Road Site. The owners of both these sites spend a great deal of time in the outdoors, either at the stables or working with equipment, and children play outdoors on the New Vernon Road Site. The activities of the residents and their customers may lead to suspended asbestos particulates in the air, thereby creating a better opportunity for exposure.

Asbestos contamination can be brought into the home as well. Studies indicate that asbestos workers have carried contamination home on their clothing and on their person. [3] In the situation at these sites, it is possible that children and pets which frequent the site may also carry contamination into the home.

The ATSDR considers the high concentration of asbestos in the soil outside the homes on the site to represent a serious hazard to the occupants' and the general public's health. Continued exposure to free asbestos fibers at concentrations present at these sites represents an imminent and substantial health hazard to exposed individuals.

#### CONCLUSIONS

Residents in homes at this site with free asbestos fibers face an imminent and substantial health hazard from exposure to asbestos. Workers at the New Vernon Road Site and workers and customers at the horse farm and riding stables at the White Bridge Road Site may also encounter asbestos contamination and face an increased risk of developing adverse health effects. The families of these workers and customers will be at risk if the free asbestos fibers are taken into their residential environments.

### RECOMMENDATIONS AND PROPOSED ACTIONS

The ATSDR will consult with the New Jersey State Department of Health on actions needed to address asbestos contamination that is not related to the NPL Site known as the "Asbestos Disposal Sites." The ATSDR, in consultation with the New Jersey State Department of Health, will develop exposure-based criteria to identify residents at risk of adverse health effects associated with these subsites. The two agencies will make a medical monitoring program available to those residents identified. A health education program for the community will be made available as well.

In addition, ATSDR recommends the following actions be taken to mitigate the health hazard associated with asbestos contamination at the New Vernon Road Site and the White Bridge Road Site:

- The EPA should dissociate affected residents, either on-site or off-site, from exposure to the site-related asbestos fibers in indoor air.
- 2. Additional sampling should be performed by the EPA to determine the extent of off-site migration.

- 3. Additional sampling for the presence of asbestos should be performed by EPA to determine if workers and customers of the affected businesses are being exposed. Initially, this sampling should be targeted at areas frequented by those workers and customers who are physically on-site for at least 40 hours per week. The sampling should include their homes. The targeting is recommended due to the longer exposures of these individuals. Additional sampling of individuals with shorter exposures may become necessary based on an evaluation of these initial results.
- 4. If rural New Jersey background levels of asbestos are not already available from State agencies, a concurrent sample to those already recommended should be collected in a maximum of three homes of similar construction in a rural setting in New Jersey. The ATSDR will accept these control samples as indicative of rural background asbestos concentrations in that part of New Jersey.
- 5. The EPA or the property owners should restrict access, authorized or unauthorized, to those areas known or suspected to be contaminated with asbestos. This restriction applies to workers, residents, and customers.
- 6. The EPA or the property owners should reduce or eliminate activities that would increase airborne particulates in those areas known or suspected to be contaminated with asbestos.
- 7. If Recommendations 5 and 6 cannot be implemented, EPA should post warning signs in the vicinity of the horse track at the White Bridge Road Site to advise customers of the asbestos-related hazards at the site.

For additional information, please contact ATSDR at the following address:

> Robert C. Williams, P.E. Director, Division of Health Assessment and Consultation Agency for Toxic Substances and Disease Registry 1600 Clifton Road, NE, MS E-32 Atlanta, Georgia 30333 (404) 639-0610 FTS 236-0610

> > 6

### REFERENCES

- 1. <u>Realth Assessment for Asbestos Disposal Site</u>. Prepared by Agency for Toxic Substances and Disease Registry. April 10, 1989.
- Revised Recommended Asbestos Standard. Prepared by National Institute for Occupational Safety and Health. December 1976.
- 3. <u>Draft Toxicological Profile for Asbestos</u>. Prepared by Agency for Toxic Substances and Disease Registry. February 16, 1990.
- 4. Excerpts from Sampling Report: New Vernon Road Site and White Bridge Road Site, Asbestos Dump Sites. Prepared by Fred C. Hart Associates, Inc., Subcontractor to Roy F. Weston, Inc. Prepared for: U.S. Environmental Protection Agency, Region II. Undated.
- 5. Conference call between Agency for Toxic Substances and Disease Registry. Division of Health Assessment and Consultation, Emergency Response and Consultation Branch (ERCB); Public Health Advisor, Agency for Toxic Substances and Disease Registry, Region II; and, New Jersey Department of Health. August 21, 1990. Refer to ERCB Superfund Record of Communication dated August 27, 1990.
- 6. Conversation between Mr. Richard Nickle, Agency for Toxic Substances and Disease Registry and Mr. William Howard, Centers for Disease Control. August 30, 1990. Refer to ATSDR Superfund Record of Communication dated August 31, 1990.
- 7. Conversation between Mr. Richard Nickle, Agency for Toxic Substances and Disease Registry and Mr. Raymond McQueen, National Asbestos Council. August 30, 1990. Refer to ATSDR Superfund Record of Communications dated August 31, 1990.
- 8. Facsimile transmission of preliminary analytical data.
  Transmitted by: Mr. Arthur Block, Public Health Advisor, ATSDR
  Regional Services, Region II, Agency for Toxic Substances and
  Disease Registry, New York. Transmitted to: Mr. Richard Nickle,
  Division of Health Assessment and Consultation, Emergency Response
  and Consultation Branch, Agency to Toxic Substances and Disease
  Registry. August 5, 1990.
- 9. Fifth Annual Report on Carcinogens: Summary 1989 (NTP 89-239).
  Prepared for the National Institute of Environmental Health
  Sciences, U.S. Public Health Service, U.S. Department of Health
  and Human Services. Prepared by: Technical Resources, Inc.,
  Rockville, Maryland.
- 10. Telephone conversations between Mr. Arthur Block, Ms. Nicki DiForte, and Mr. Richard Nickle concerning air sampling results. Conversation 10/24 and 10/25, 1990. Referred to ATSDR Superfund Record of Communications dated 10/25/90.

11. New Vernon Residential Asbestos Air Investigation: Meyersville,
New Jersey. Prepared for the EPA/U.S. EPA Environmental Response
Team (ERT). Prepared by ERT's REAC contractor, Edison, New
Jersey. No date.

ABD 001 2116